

NBI-HE-92-43

July 1992

Free Field Representations and Screening Operators for the $N = 4$ Doubly Extended Superconformal Algebras

Katsushi Ito, Jens Ole Madsen and Jens Lyng Petersen

*The Niels Bohr Institute, Blegdamsvej 17
DK-2100, Copenhagen Ø, Denmark*

Abstract

We present explicit free field representations for the $N = 4$ doubly extended superconformal algebra, $\tilde{\mathcal{A}}_\gamma$. This algebra generalizes and contains all previous $N = 4$ superconformal algebras. We have found $\tilde{\mathcal{A}}_\gamma$ to be obtained by hamiltonian reduction of the Lie superalgebra $D(2|1;\alpha)$. In addition, screening operators are explicitly given and the associated singular vectors identified. We use this to present a natural conjecture for the Kac determinant generalizing a previous conjecture by Kent and Riggs for the singly extended case. The results support and illuminate several aspects of the characters of this algebra previously obtained by Taormina and one of us.

In recent works, we [1] and two of us [2], have outlined how essentially all known superconformal algebras – including (super- W -type) ones (for $N \geq 4$) for which the algebra only closes on composites of generators – may be classified and indeed be given a unified treatment, using the techniques of hamiltonian reductions of WZNW models on Lie (super)groups [3].

In ref.[2] this program was described at the level of the *classical* hamiltonian reduction, whereas in ref. [1] the quantum case was considered as well using explicit free field realizations. In the case of superconformal algebras with an affine $\widehat{SO}(N)$ algebra such a realization has been known for some time [4, 5]. In [1] it was further discussed and explicit realizations were described for the non-exceptional type Lie superalgebras, $sl(N|2)$, $osp(N|2)$ and $osp(4|2N)$. For the $N = 4$ singly extended algebra containing an affine $\widehat{SU}(2)$ [6] it has been recently provided by Matsuda, [7]. In the present letter we describe the case of the $N = 4$ “doubly extended” superconformal algebra of Sevrin, Troost and Van Proeyen [8] (see also [9]-[11]), commonly denoted $\tilde{\mathcal{A}}_\gamma$ (see [12]-[14]) and further analyzed by several authors ([15]-[17]). The algebra, $\tilde{\mathcal{A}}_\gamma$, was originally given in a form with auxiliary free fermionic generators and an extra $U(1)$ current. That algebra is denoted $\mathcal{A}_\gamma$ and has the superalgebra $D(2|1; \alpha)$ as its finite dimensional sub-algebra. We have found that $\tilde{\mathcal{A}}_\gamma$ is obtained as a result of carrying out the hamiltonian reduction of the WZNW model on the Lie supergroup of the Lie superalgebra $D(2|1; \alpha)$.

We first provide a brief account of how to translate between the notations of [1] using the Lie superalgebra language, and the original notations for $\tilde{\mathcal{A}}_\gamma$ (see for example [8, 12]).

The exceptional Lie superalgebra, $D(2|1; \alpha)$ (cf. ref. [18]) depends on the parameter α or $\gamma \equiv \frac{\alpha}{1-\alpha}$, and has rank three. Define the simple roots, $\alpha_1, \alpha_2, \alpha_3$ with metric given by

$$\begin{aligned} \alpha_1^2 &= 0, & \alpha_2^2 &= -2\gamma, & \alpha_3^2 &= -2(1 - \gamma), \\ \alpha_1 \cdot \alpha_2 &= \gamma, & \alpha_1 \cdot \alpha_3 &= 1 - \gamma, & \alpha_2 \cdot \alpha_3 &= 0, \end{aligned} \tag{1}$$

Then the positive (mutually orthogonal) even roots are α_2, α_3 and $\theta \equiv 2\alpha_1 + \alpha_2 + \alpha_3 \equiv \alpha_{1123}$, with $\theta^2 = 2$. They define the even subalgebra, $\mathfrak{g}_0 = A_1 \oplus A_1 \oplus A_1$, or (partly in the notation of [8]), $su(2)^+ \oplus su(2)^- \oplus su(2)_\theta$. α_2, α_3 define the set $\Delta_+^0(G)$ with $G = su(2)^+ \oplus su(2)^-$ in the notation of [1]. It is the affine $\widehat{SU}(2)_\theta$ algebra defined by θ which is constrained in the hamiltonian reduction and gives rise to the Virasoro subalgebra.

The positive odd roots forming the set Δ_+^1 are given by

$$\begin{aligned} \alpha_1 &\equiv \beta_-, & \alpha_{12} &\equiv \alpha_1 + \alpha_2 \equiv \beta_{+K}, \\ \alpha_{13} &\equiv \alpha_1 + \alpha_3 \equiv \beta_{-K}, & \alpha_{123} &\equiv \alpha_1 + \alpha_2 + \alpha_3 \equiv \beta_+, \end{aligned} \quad (2)$$

the $+, -, +K, -K$ notation being used in the original literature on $\tilde{\mathcal{A}}_\gamma$. The pairing $+\leftrightarrow -$ and $+K \leftrightarrow -K$ corresponds to the pairing $\beta \leftrightarrow \theta - \beta$ (with $\beta, \theta - \beta \in \Delta_+^1$) in the notation of [1]. The vector space, $\mathfrak{g}_1$, of the odd generators carry a representation of $G = su(2)^+ \oplus su(2)^-$ the representation matrices of which may be obtained as follows:

For $\beta, \beta' \in \Delta_+^1$, and $\alpha \equiv \beta' - \beta \in \Delta^0$, define

$$t_{\theta-\beta',\beta}^\alpha \equiv 2 \frac{N_{-\beta',\beta}}{\alpha^2} = N_{\alpha,\beta} \quad (3)$$

where $N_{\beta',\beta}$ are the structure coefficients of (the odd-odd part of) $D(2|1;\alpha)$. The relation to the notation of [8] consists in denoting

$$+\alpha_2 : (++) , \quad -\alpha_2 : (+-), \quad +\alpha_3 : (-+), \quad -\alpha_3 : (--)$$

One then finds

$$t_{+,+K}^{+-} = +1, \quad t_{-,-K}^{++} = -1, \quad t_{-,+K}^{-+} = -1, \quad t_{+,-K}^{--} = +1, \quad (4)$$

the remaining ones being zero. Matrix multiplication is defined by

$$(t^\alpha t^{\alpha'})_{\theta-\beta',\beta} \equiv \sum_{\beta'' \in \Delta_+^1} t_{\theta-\beta',\beta''}^\alpha t_{\theta-\beta'',\beta}^{\alpha'}. \quad (5)$$

Define t^2 and t^3 (corresponding to $2t^{+3}$ and $2t^{-3}$ in the traditional notation) by $[t^{\alpha_i}, t^{-\alpha_i}] = t^i$. Then

$$\pm t_{+K,-K}^{\pm 3} = \mp t_{-K,+K}^{\pm 3} = \frac{1}{2}, \quad t_{+,-}^{\pm 3} = -t_{-,+}^{\pm 3} = \frac{1}{2}. \quad (6)$$

Thus we have the two commuting doublet representations of the two $SU(2)$'s ($i = 2, 3$):

$$\vec{t}_i \equiv (t^{\alpha_i}, t^{-\alpha_i}, t^i) = (t^{\pm+}, t^{\pm-}, 2t^{\pm 3}) \equiv \vec{t}^{\pm}.$$

The space, $\mathfrak{g}_1 = \mathfrak{g}_1^+ \oplus \mathfrak{g}_1^-$ falls in two parts corresponding to positive and negative odd roots (with isospins $\pm \frac{1}{2}$ according to $su(2)_\theta$, cf. [1]). Any basis for $\mathfrak{g}_1^+$ is indexed by the positive odd roots $\beta \in \Delta_+^1$. We denote the $su(2)^+ \oplus su(2)^-$ transformation of such a generic element X_β , by

$$t(X_\beta) = (tX)_\beta = \sum_{\beta' \in \Delta_+^1} t_{\beta, \beta'} X_{\theta - \beta'} \quad (7)$$

We now turn to a brief description of the classical superconformal algebra, $\tilde{\mathcal{A}}_\gamma^{cl}$, to be compared below with the well known quantum $\tilde{\mathcal{A}}_\gamma$, [12, 13, 14, 8]:

Besides the energy-momentum tensor, $L(z)$, with (classical) central charge $c_{cl} = -6k$, it has the following generators: Two sets of commuting affine $\widehat{SU}(2)^\pm$ currents, related to the currents of the (classical) even subalgebras of the affine Lie superalgebra $\{J_{\alpha_i}, H^i\}$, with $\alpha_i \in \Delta(G)$, for $i = 2, 3$ with (classical) level, k .¹

Denote $H_{\alpha_i} \equiv \frac{2\alpha_i \cdot H}{\alpha_i^2}$ and write

$$\vec{J}_i \equiv (J_{\alpha_i}, J_{-\alpha_i}, H_{\alpha_i}) = (T^{\pm+}, T^{\pm-}, 2T^{\pm 3}) \equiv \vec{J}^{\pm},$$

for $i = 2, 3$. Introduce $\vec{t}_i \cdot \vec{J}_i \equiv \frac{1}{2}[t^{\alpha_i} J_{-\alpha_i} + t^{-\alpha_i} J_{\alpha_i} + t^i H_{\alpha_i}]$ and $\vec{J}_i^2 \equiv \frac{1}{2}[J_{\alpha_i} J_{-\alpha_i} + J_{-\alpha_i} J_{\alpha_i} + H_{\alpha_i}^2]$, (normal ordering of modes implied). These affine currents satisfy

$$\begin{aligned} J_{\alpha_i}(z) J_{-\alpha_i}(w) &= \frac{\frac{2k}{\alpha_i^2}}{(z-w)^2} + \frac{H_{\alpha_i}(w)}{z-w} + \dots, \\ H_{\alpha_i}(z) H_{\alpha_i}(w) &= \frac{\frac{4k}{\alpha_i^2}}{(z-w)^2} + \dots, \quad H_{\alpha_i}(z) J_{\pm \alpha_i}(w) = \pm \frac{2J_{\pm \alpha_i}(w)}{z-w} + \dots, \end{aligned} \quad (8)$$

¹Notice that k is negative for a unitary representation in our notation since we are using negative definite metric for the even roots. We change to a more standard notation presently.

and they are conformal primaries with dimension 1. All these OPE's are understood here in the classical case in the formal sense of the hamiltonian structure.

Next there are four supercurrents, $G_\beta(z)$ indexed by the positive odd roots, related to the notation $(+, -, +K, -K)$ of [8, 12] as above. They are conformal primaries with dimension, $3/2$, and affine primaries transforming as $(i = 2, 3)$

$$J_{\alpha_i}(z)G_\beta(w) = \frac{(t^{\alpha_i}G)_\beta(w)}{z-w} + \dots, \quad H_{\alpha_i}(z)G_\beta(w) = \frac{(t^i G)_\beta(w)}{z-w} + \dots \quad (9)$$

The general result of [1] for the classical OPE's of the supercurrents reduces in this case to

$$\begin{aligned} G_{\theta-\beta'}(z)G_\beta(w) &= -2k\delta_{\beta,\beta'}(z-w)^{-3} + 2(z-w)^{-2}M_{\theta-\beta',\beta}(w) + (z-w)^{-1}\partial_w M_{\theta-\beta',\beta}(w) \\ &+ \delta_{\beta,\beta'}(z-w)^{-1} \left[L(w) + \frac{1}{K} \{ \vec{J}_2^2 + \vec{J}_3^2 + t_{\theta-\beta,\beta}^2 H_{\alpha_2}(w) H_{\alpha_3}(w) \} \right] + \dots, \end{aligned} \quad (10)$$

where

$$M_{\theta-\beta',\beta} \equiv \left[\gamma \vec{t}_2 \cdot \vec{J}_2 + (1-\gamma) \vec{t}_3 \cdot \vec{J}_3 \right]_{\theta-\beta',\beta}. \quad (11)$$

It is convenient to reparametrize the two constants, k, γ in terms of $K^\pm$ (and $K \equiv K^+ + K^-$) thus

$$\gamma = \frac{K^-}{K}, \quad 1-\gamma \equiv \frac{K^+}{K}, \quad k = -\frac{K^+K^-}{K}, \quad (12)$$

showing that the parameters, $K^\pm$, are simply the classical levels of the two commuting, affine $\widehat{SU}(2)$'s.

The *quantum* algebra, $\tilde{\mathcal{A}}_\gamma$, has an entirely similar form with the OPE's having the standard meaning in conformal field theory. However, some of the parameters get “renormalized”. Thus the energy momentum tensor has central charge [12, 1]

$$c = -6k - 3 = 6\frac{K^+K^-}{K} - 3. \quad (13)$$

The two commuting affine $\widehat{SU}(2)$'s have levels $\tilde{K}^\pm$ rather than $K^\pm$, where $\tilde{K}^\pm \equiv K^\pm - 1$. Similarly, the operator, $M_{\theta-\beta',\beta}$ is modified into

$$M_{\theta-\beta',\beta} = \left[\frac{\tilde{K}^-}{K} \vec{t}^+ \cdot \vec{J}^+ + \frac{\tilde{K}^+}{K} \vec{t}^- \cdot \vec{J}^- \right]_{\theta-\beta',\beta}. \quad (14)$$

We now turn to the explicit free field representation of this quantum algebra, cf. [1]. We introduce commuting sets of affine currents ($i = 2, 3$),

$$\vec{J}_i \equiv (\hat{J}_{\alpha_i}, \hat{J}_{-\alpha_i}, \hat{H}_{\alpha_i}), \quad (15)$$

with similar OPE's as the currents $\vec{J}_i$, except that they correspond to Kac-Moody levels, $\tilde{K}^\pm - 1$ rather than $\tilde{K}^\pm$. We could introduce the standard Wakimoto free field representations [19] for those as well, but we shall not need these explicitly. We further introduce free fermions, $\chi_\beta(z)$, with OPE's

$$\chi_{\theta-\beta'}(z)\chi_\beta(w) = \frac{\delta_{\beta'\beta}}{z-w} + \dots \quad (16)$$

They are taken to commute with the currents, $\vec{J}_i(z)$, $i = 2, 3$. Finally a free scalar, $\phi(z)$ coupled to world sheet curvature is used. Based on these we first introduce fermionic affine $\widehat{SU}(2)_1 \times \widehat{SU}(2)_1$ currents

$$\vec{J}_{F,i}(z) \equiv \sum_{\beta, \beta' \in \Delta_+^1} \chi_{\theta-\beta}(z)(\vec{t}_{\beta, \theta-\beta'})_i \chi_{\beta'}(z). \quad (17)$$

Following ref. [1] we write (the dual Coxeter number, $h^\vee$ of the exceptional Lie superalgebra, $D(2|1; \alpha)$, is zero)

$$\begin{aligned} \alpha_+ &\equiv \sqrt{k + h^\vee} = -i\sqrt{\frac{K^+ K^-}{K}}, \\ Q &\equiv \sqrt{\theta^2} \left(\frac{2 - |\Delta_+^1|}{2\alpha_+} - \frac{2\alpha_+}{\theta^2} \right) = -\sqrt{2}(\alpha_+^{-1} + \alpha_+). \end{aligned} \quad (18)$$

Also

$$L_\phi = -\frac{1}{2}(\partial\phi)^2 - \frac{iQ}{2}\partial^2\phi, \quad L_F = \frac{1}{2} \sum_{\beta \in \Delta_+^1} \partial\chi_{\theta-\beta}\chi_\beta, \quad L_{J^\pm} = \frac{2}{K^\pm}(\vec{J}^\pm)^2. \quad (19)$$

It is then a straightforward (albeit tedious) task to check that the following provides the sought free field realization of the algebra:

$$\begin{aligned} L(z) &= L_{J^+}(z) + L_{J^-}(z) + L_F(z) + L_\phi(z), \\ \vec{J}^\pm(z) &= \vec{J}^\pm(z) + \vec{J}_F^\pm(z), \\ G_\beta(z) &= \left\{ \frac{Q}{\sqrt{2}}\partial + \frac{i}{\sqrt{2}}\partial\phi + \frac{2\gamma}{\alpha_+}(\vec{J}^+ + \frac{3}{2}\vec{J}_F^+) \cdot \vec{t}^+ + \frac{2(1-\gamma)}{\alpha_+}(\vec{J}^- + \frac{3}{2}\vec{J}_F^-) \cdot \vec{t}^- \right\} \chi_\beta(z). \end{aligned} \quad (20)$$

It is easy to verify that these expressions generalize the ones previously presented in the literature. Thus, for $\gamma = \frac{1}{2}$ or $K^+ = K^-$, the algebra $\tilde{\mathcal{A}}_\gamma$ reduces to the $N = 4$ superconformal algebra with an affine $\widehat{SO}(4)$, and the above expressions to the ones given by Mathieu [4]. In particular one may easily verify that $\vec{J}_F^+ \cdot \vec{t}^+ \chi_\beta(z) \equiv -\vec{J}_F^- \cdot \vec{t}^- \chi_\beta(z)$ so that these (triple fermionic) terms cancel in that limit.

Second, in the limit $\gamma \rightarrow 1$, or $K^- \rightarrow \infty$, or $\alpha_+ \rightarrow -i\sqrt{K^+}$ (with a similar “charge exchanged” limit), the $\tilde{\mathcal{A}}_\gamma$ algebra reduces to the singly extended $N = 4$ algebra with a single affine $\widehat{SU}(2)$. In that limit the above expressions reduce to the ones recently given by Matsuda [7].

These results have immediate consequences for the representation theory of the $\tilde{\mathcal{A}}_\gamma$ algebra, as studied in [12, 15, 16]. In particular, it follows that characters of highest weight representations (for details of the definition of $\tilde{\mathcal{A}}_\gamma$ characters, see [15]) take the form

$$Ch^{\tilde{\mathcal{A}}_\gamma}(h, K^+, K^-, \ell^+, \ell^-; q, z_+, z_-) = \chi_\phi(q) \chi_F(q, z_+, z_-) \chi_{2\ell^+}^{\tilde{K}^+-1}(q, z_+) \chi_{2\ell^-}^{\tilde{K}^--1}(q, z_-) \quad (21)$$

where

$$\chi_\phi(q) = \frac{q^{h_\phi + \frac{Q^2}{8}}}{\eta(q)} \quad (22)$$

is the character of the ϕ -module with $c_\phi = 1 - 3Q^2$ and conformal dimension, h_ϕ .

$$\chi_F(q, \xi_+, \xi_-) = \frac{\theta_3(q, z_+) \theta_3(q, z_-)}{\eta^2(q)} \quad (23)$$

is the contribution from the 4 fermions. For simplicity we only give this expressions in the NS sector, but the translation to the R sector is trivial using standard methods, cf. [17]. Finally $\chi_{2\ell}^k(q, z)$ is the standard affine $\widehat{SU}(2)_k$ character for isospin ℓ and level k . This expression for the character is in complete agreement with what was found in ref. [15] for the characters of unitary, irreducible, *massive* representations. In [15] it looked miraculous (i) that the factorized form should appear, and (ii) that affine $\widehat{SU}(2)$ characters at levels $\tilde{K}^\pm - 1$ should occur. These aspects now are fully explained.

As shown in [12] unitary representations of $\tilde{\mathcal{A}}_\gamma$ occur only when (a) we have integrable representations of affine $\widehat{SU}(2)_{\tilde{K}^+} \times \widehat{SU}(2)_{\tilde{K}^-}$ (i.e.: $K^\pm$ are positive integers, $2\ell^\pm$ are positive integers $\leq \tilde{K}^\pm$ in the NS sector), and (b) when the conformal dimension in the NS sector is bounded by

$$h \geq h_0^{NS} \equiv \frac{1}{K} \{(\ell^+ - \ell^-)^2 + K^+ \ell^- + K^- \ell^+\}. \quad (24)$$

The corresponding bound in the R sector is given by

$$h \geq h_0^R \equiv \frac{1}{K} (\ell^+ + \ell^-)^2 + \frac{K^+ K^-}{4K} - \frac{1}{4}. \quad (25)$$

At the limiting value, the massive representation is no longer irreducible, but splits into two massless or chiral representations.

Clearly, for $h_\phi = -\frac{Q^2}{8}$ (which is positive for integrable representations) the characters transform according to a finite dimensional representation of the modular group.

We now turn to the determination of screening operators relevant for the discussion of null states in degenerate representations. Again, these will turn out to be closely related to but generalizing the known expressions [4, 7] for the two standard versions of $N = 4$ superconformal algebras. Screening operators associated with the affine $\widehat{SU}(2)_{\tilde{K}^+ - 1} \times \widehat{SU}(2)_{\tilde{K}^- - 1}$ are standard [19], and will not be further discussed here (cf. also [1]). In addition to those we find two other screening operators (as in [1]).

We first introduce the primary multiplet of fields,

$$\Phi_{\vec{\lambda}}^{\overline{\Lambda}^*}(z) \equiv [\Phi_{m^+}^{\frac{1}{2}}(z)]^{(+)} [\Phi_{m^-}^{\frac{1}{2}}(z)]^{(-)}$$

of the currents $\vec{J}_i$, carrying the $\overline{\Lambda}^* = (\frac{1}{2}, \frac{1}{2})$ representation of $SU(2)^+ \times SU(2)^-$, with weight $\overline{\Lambda} = (m^+, m^-) = (\pm\frac{1}{2}, \pm\frac{1}{2})$. These weights may naturally be identified with the positive, odd roots, and phases may be chosen so that

$$t^\alpha(\Phi_{\vec{\beta}}^{\overline{\Lambda}^*}(z)) = \Phi_{\vec{\beta}+\vec{\alpha}}^{\overline{\Lambda}^*}(z) \quad (26)$$

for $\alpha \in \Delta(G)$ and $\beta, \beta + \alpha \in \Delta_+^1$. Then the fermionic screening operator may be given as

$$S_f(z) = \sum_{\beta \in \Delta_+^1} \chi_\beta(z) \Phi_{\theta-\beta}^{\bar{\Lambda}^*}(z) \exp\left(-\frac{i\sqrt{\theta^2}\phi(z)}{2\alpha_+}\right), \quad (27)$$

in accord with the structure of the general formula in [1]. A detailed calculation shows that

$$G_\beta(z)S_f(w) = \partial_w \left(\frac{\Phi_\beta^{\bar{\Lambda}^*}(w) \exp\left(-\frac{i\sqrt{\theta^2}\phi(w)}{2\alpha_+}\right)}{z-w} \right) + \dots \quad (28)$$

In comparison with the result in ref.[7] it may be noticed that the sum over β involves 4 terms which pairwise form singlets of either $SU(2)^+$ or $SU(2)^-$ but not of both. In the limit $K^- \rightarrow \infty$ (say) where one of these affine algebras are frozen out, it is only important to have singlets of $\widehat{SU}(2)_{\bar{K}^+}^+$ so that one actually finds two independent fermionic screening operators in that limit as in ref. [7]. In general, however, there is only the single one we have provided. Nevertheless the same number of families of null states are generated by our single chiral screening operator as with the two of [7], cf. the discussion of the Kac determinant below.

As argued in [1] we expect another screening operator of the form

$$S_\theta(z) = s_\theta(z) \exp\left(\frac{2i\alpha_+\phi(z)}{\sqrt{\theta^2}}\right), \quad (29)$$

in order to characterize the singular vectors for the boson ϕ . The explicit form for $s_\theta(z)$ has been obtained previously only in a limited number of cases. In the present one we find it may be expressed (for example) as follows.

$$\begin{aligned} s_\theta(z) &= (\alpha_+^2 + 1)\chi_+(z)\chi_-(z)\chi_{+K}(z)\chi_{-K}(z) + (1 - 2\gamma)[L_\phi(z) - \frac{i\alpha_+}{\sqrt{2}}\partial^2\phi(z) - L_F(z)] \\ &- \gamma(L_{\hat{j}^+} - 2\vec{\hat{J}}^+ \cdot \vec{J}_F^+)(z) + (1 - \gamma)(L_{\hat{j}^-} - 2\vec{\hat{J}}^- \cdot \vec{J}_F^-)(z) \end{aligned} \quad (30)$$

again this expression reduces to the corresponding ones in the limits, $\gamma \rightarrow \frac{1}{2}, 1$, [4, 7], and for the (“most non-trivial”) OPE’s one finds

$$\begin{aligned} G_\beta(z)S_\theta(w) &= \partial_w \left\{ (z-w)^{-1} \left\{ [i\alpha_+(2\gamma-1)(\partial + \frac{1}{\sqrt{2}}\partial\phi) + 2\gamma(\vec{\hat{J}}^+ \cdot \vec{t}^+ - \frac{2}{3}(\alpha_+^2 + 1)\vec{J}_F^+ \cdot \vec{t}^+)] \right. \right. \\ &- \left. \left. 2(1-\gamma)(\vec{\hat{J}}^- \cdot \vec{t}^- - \frac{2}{3}(\alpha_+^2 + 1)\vec{J}_F^- \cdot \vec{t}^-) \right] \chi_\beta(w) \right\} e^{\frac{2i\alpha_+\phi(w)}{\sqrt{\theta^2}}} + \dots \end{aligned} \quad (31)$$

The singular vectors produced by integrating the screening charges for the affine $\widehat{SU}(2)_{\tilde{K}+1} \times \widehat{SU}(2)_{\tilde{K}-1}$ (which we did not explicitly provide) may rather easily be seen to produce (i) for integrable representations, precisely the singular vectors constructed and used in ref.[15] for building characters for the unitary representations of the $\tilde{\mathcal{A}}_\gamma$ algebra (generalizing the techniques of [20, 21]), and (ii) singular vectors associated with non-unitary, non-integrable representations in complete analogy to the case of the singly extended $N = 4$ theory, cf. [7]. In addition to those, there are singular vectors produced by the screening operators, $S_f(z)$ and $S_\theta(z)$, just given. Again the situation is rather analogous to the one discussed in the special case $\gamma \rightarrow 1$ by Matsuda [7]. In that case families of null states were conjectured based on computer studies by Kent and Riggs [22] some time ago. In [7] it was shown how these were exactly produced by the corresponding screening operators. Consistency of the characters demands that these extra singular vectors lie entirely outside the unitarity domain². Further, once the multiplicity of the zeros in the Kac-determinant corresponding to all these singular vectors are understood and found to saturate the bound provided by the corresponding level, one may consider that the final part of a rigorous proof has been provided for the characters, [20, 15], and the Kac determinant as well.

In the present letter we confine ourselves to presenting a natural proposal for how the Kent and Riggs conjecture for the Kac determinant should generalize for a general value of the parameter, $\gamma \neq 1$. For simplicity we consider only the R-sector as in [7]. It is straight forward to repeat the treatment in the NS sector or in fact to use “spectral flow” from the R sector result given here, [17]. We attempt a notation making the comparison with ref. [22] easy.

The structure of highest weight multiplets in the R sector is as follows (for a complete discussion, cf. [15]): The massive representation has 4 $SU(2)^+ \times SU(2)^-$ multiplets with

²One of us (JLP) is indebted to A. Kent for illuminating discussions.

the same (lowest) value of the conformal dimension, h , and isospins parametrized as

$$(\ell^+, \ell^- - \frac{1}{2}), (\ell^+ - \frac{1}{2}, \ell^-), (\ell^+ - 1, \ell^- - \frac{1}{2}), (\ell^+ - \frac{1}{2}, \ell^- - 1).$$

The representation is referred to by means of the triple (h, ℓ^+, ℓ^-) in addition to the two parameters, (k, γ) , or equivalently, (K^+, K^-) . The 4 isospin multiplets are connected by the action of the 4 supercurrent zero-modes. The two first members above are annihilated by $(G_+)_0, (G_{+K})_0$ and $(G_+)_0, (G_{-K})_0$, respectively. The Kac determinant is the determinant of the sesquilinear form of inner products of states in the module built on the highest weight multiplet, and having definite values of the quantum numbers, thus h, ℓ_3^+, ℓ_3^- are increased respectively by $n, \Delta\ell^+, \Delta\ell^-$. It is denoted by $M_{n, \Delta\ell^+, \Delta\ell^-}(h, k, \gamma, \ell^+, \ell^-)$.

Multiplicities of states in this subspace are given by partition functions, $P_\gamma^4(n, 2\Delta\ell^+, 2\Delta\ell^-)$ which are simple to describe provided all generators act freely. In that case they are generated as follows

$$\begin{aligned} P_\gamma^4(q, z_+, z_-) &\equiv \prod_{n=1}^{\infty} \frac{(1 + q^n z_+ z_-)(1 + q^n z_+ z_-^{-1})(1 + q^n z_+^{-1} z_-)(1 + q^n z_+^{-1} z_-^{-1})}{(1 - q^3)(1 - q^n z_+^2)(1 - q^n z_+^{-2})(1 - q^n z_-^2)(1 - q^n z_-^{-2})} \\ &\times \frac{(1 + z_+^{-1} z_-)(1 + z_+^{-1} z_-^{-1})}{(1 - z_+^{-2})(1 - z_-^{-2})} \\ &\equiv \sum_{n, 2\Delta\ell^+, 2\Delta\ell^-=1}^{\infty} q^n z_+^{2\Delta\ell^+} z_-^{2\Delta\ell^-} P_\gamma^4(n, 2\Delta\ell^+, 2\Delta\ell^-). \end{aligned} \quad (32)$$

In case the supercurrent with $SU(2)^+ \times SU(2)^-$ weight $(\frac{1}{2}\epsilon^+, \frac{1}{2}\epsilon^-)$ and level, N does *not* act freely (i.e. is expressible in terms of other operators), the multiplicity is provided by the modified functions

$$\begin{aligned} P_\gamma^4(q, z_+, z_-; N, \epsilon^+, \epsilon^-) &\equiv P_\gamma^4(q, z_+, z_-)(1 + q^N z_+^{\epsilon^+} z_-^{\epsilon^-})^{-1} \\ &\equiv \sum_{n, 2\Delta\ell^+, 2\Delta\ell^-=1}^{\infty} q^n z_+^{2\Delta\ell^+} z_-^{2\Delta\ell^-} P_\gamma^4(n, 2\Delta\ell^+, 2\Delta\ell^-; N, \epsilon^+, \epsilon^-). \end{aligned} \quad (33)$$

We further generalize the various functions of [22] (corresponding to $\rho = \eta = 0$ in their notation; this is the R sector), the vanishing of which corresponds to the occurrence of

the singular states as described below. Thus we define

$$\begin{aligned}
f_{p,q}(h, k, \gamma, \ell^+, \ell^-) &= k(k+1+4h) - (kp-q)^2 + 4(\gamma(\ell^+)^2 + (1-\gamma)(\ell^-)^2), \\
g_{p^\pm, q^\pm}(h, K^\pm, \ell^\pm) &= K^\pm |p^\pm| + 2\text{sgn}(p^\pm)\ell^\pm - q^\pm, \\
h_{N, \epsilon^+, \epsilon^-}(h, k, \gamma, \ell^+, \ell^-) &= 4h - k(4N^2 - 1) + 1 + 8N(\gamma\epsilon^+\ell^+ + (1-\gamma)\epsilon^-\ell^-) \\
&\quad - \frac{4}{K}(\epsilon^+\ell^+ - \epsilon^-\ell^-)^2.
\end{aligned} \tag{34}$$

These are simply related to corresponding functions in [22] in the limit $\gamma \rightarrow 1$. The precise form follows from the identification of singular states presented below.

The corresponding conjecture for the Kac determinant takes the form

$$\begin{aligned}
M_{n, \Delta\ell^+, \Delta\ell^-}(h, k, \gamma, \ell^+, \ell^-) &= A \prod_{p, q > 0} [f_{p,q}(h, k, \gamma, \ell^+, \ell^-)]^{P_\gamma^4(n-pq, 2\Delta\ell^+, 2\Delta\ell^-)} \\
&\times \prod_{\substack{p^+ = -\infty \\ q^+ = 1}}^{\infty} [g_{p^+, q^+}(h, K^+, \ell^+)]^{P_\gamma^4(n-|p^+|q^+, 2\Delta\ell^+ + \text{sgn}(p^+)q^+, 2\Delta\ell^-)} \\
&\times \prod_{\substack{p^- = -\infty \\ q^- = 1}}^{\infty} [g_{p^-, q^-}(h, K^-, \ell^-)]^{P_\gamma^4(n-|p^-|q^-, 2\Delta\ell^+, 2\Delta\ell^- + \text{sgn}(p^-)q^-)} \\
&\times \prod_{\substack{N=1 \\ \epsilon^\pm = \pm 1}}^{\infty} [h_{N, \epsilon^+, \epsilon^-}(h, k, \gamma, \ell^+, \ell^-)]^{P_\gamma^4(n-N, 2\Delta\ell^+ - \epsilon^+, 2\Delta\ell^- - \epsilon^-; N, \epsilon^+, \epsilon^-)} \\
&\times \prod_{\epsilon^- = \pm 1} [h_{0,1, \epsilon^-}(h, k, \gamma, \ell^+, \ell^-)]^{P_\gamma^4(n, 2\Delta\ell^+ - 1, 2\Delta\ell^- - \epsilon^-; 1, \epsilon^+)} \tag{35}
\end{aligned}$$

notice that

$$\begin{aligned}
h_{0,1,-1}(h, k, \gamma, \ell^+, \ell^-) &= 4(h - \{\frac{1}{K}(\ell^+ + \ell^-)^2 + \frac{K^+ K^-}{4K} - \frac{1}{4}\}) \\
&= 4(h - h_0^R)
\end{aligned} \tag{36}$$

where (as above) h_0^R is the massless bound on the conformal dimension in the R sector. The factor A is a normalization factor.

We now briefly indicate how the screening operators we have found give rise to singular vectors which in turn give rise to the above factors in the Kac determinant.

In our free field representation a generic state is a tensor product of states pertaining to (i) the Fock space of the 4 free fermions, (ii) a representation space of the two commuting affine currents, $\vec{J}^\pm$, and (iii) the scalar, ϕ : $|F\rangle \otimes |\hat{\ell}^+\rangle \otimes |\hat{\ell}^-\rangle \otimes |p\rangle$. We refer to p as the “momentum”, the corresponding state being created by a vertex operator $e^{i\sqrt{\theta^2}p\phi(z)}$.

We first consider singular vectors produced by p -fold integrals of the ϕ -screening operator, S_θ , carrying momentum, α_+ . Consider acting with the p -fold integral on the state $|0\rangle_R \otimes |\hat{\ell}^\pm = \ell^\pm - \frac{1}{2}\rangle \otimes |P - p\alpha_+\rangle$. Here the combined Ramond vacuum states $|0\rangle_R$ of the 4 fermions provide the multiplet structure in the R sector described above. They transform as a $(\frac{1}{2}, \frac{1}{2})$ representation and provides a contribution of 4/16 to the conformal dimension. The action of the p -fold integral of S_θ will produce a singular vector which itself is a descendant of a highest weight vector with momentum, P , and isospins unchanged. The on-shell condition that the level (increase in conformal dimension) is a positive integer, $N = p \cdot q$ gives rise to the quantization of momenta

$$P = P_{p,q} = -\frac{1}{2\alpha_+}(q-1) + \frac{\alpha_+}{2}(p+1), \quad (37)$$

which is a generalization of [7] and conforms with [1] (after $P \rightarrow -Q - P$). The corresponding conformal dimension takes a value given by the vanishing of $f_{p,q}(h, k, \gamma, \ell^+, \ell^-)$ and justifies the first factor in the Kac determinant. One also checks that the corresponding values of h lie below the unitarity bound, so that these singular vectors do not affect the calculation of characters of unitary representations, [15].

The identification of singular vectors arising from the well known screening charges (not repeated here) of the affine $\widehat{SU}(2)_{\tilde{K}+1} \times \widehat{SU}(2)_{\tilde{K}-1}$ is entirely analogous to the singly extended case, [7], and gives rise to the second and third factor in the Kac determinant. For integrable representations these are exactly the ones employed in the construction of characters in ref. [15].

Singular vectors produced by the fermionic screening operator, S_f , are obtained by integrating that once only, and acting on a suitable state (cf. [23]), which we take to be

$|0\rangle_R \otimes |\hat{j}^+, \hat{j}^-\rangle \otimes |P + \frac{1}{2\alpha_+}\rangle$, since the chiral screening operator has momentum, $-1/2\alpha_+$. The result will be a singular vector with $SU(2)^+ \times SU(2)^-$ weight, $(\ell_S^+, \ell_S^-) = (\hat{j}^+ + \frac{1}{2}, \hat{j}^- + \frac{1}{2})$ (the extra $(\frac{1}{2}, \frac{1}{2})$ coming as before from the Ramond ground state multiplet of the four fermions). The singular vector is a descendant of a highest weight state with momentum, P and $SU(2)^+ \times SU(2)^-$ weight $(\ell^+, \ell^-) = (\hat{j}^+ + \frac{1}{2} + \epsilon^+ \frac{1}{2}, \hat{j}^- + \frac{1}{2} + \epsilon^- \frac{1}{2})$ provided the on-shell condition that the level is a positive integer (or zero) is fulfilled. Here there are four possibilities corresponding to $\epsilon^\pm = \pm 1$. The reason is that the screening operator, S_f involves the operators, $[\Phi_{m^\pm}^{\frac{1}{2}}]^\pm$ transforming as $(\frac{1}{2}, \frac{1}{2})$ primaries of the currents, $\vec{J}^\pm$. More explicitly the 4 possibilities may be seen by using standard free field representations for these operators. These representations involve a vertex operator or it's dual and they give rise to the total of 4 possibilities. If the level is denoted by N the quantization condition for the momentum is

$$P = P_N = \alpha_+ N + \frac{1}{2}\alpha_+ + \alpha_+ \left[\frac{\epsilon^+ \ell^+ - \frac{1}{2}}{K^+} + \frac{\epsilon^- \ell^- - \frac{1}{2}}{K^+} \right]. \quad (38)$$

This gives rise to a value for the conformal dimension provided by the vanishing of the function $h_{N, \epsilon^+, \epsilon^-}(h, k, \gamma, \ell^+, \ell^-)$ above. These conformal dimensions all lie below the unitarity bound, h_0^R , except for $N = 0, \epsilon^+ = -\epsilon^-$ which is precisely the condition for masslessness. Thus the existence of these states is also consistent with the character calculation, [15]. Finally they give rise to the remaining two factors in the Kac determinant.

The multiplicities here are given under the assumption that a certain supercurrent mode is redundant. Such phenomena are well known (see for example [20] in the singly extended $N = 4$ case, and [15] in the doubly extended $N = 4$ case) in similar contexts, but in this letter we have not in fact attempted a full understanding.

Acknowledgements:

One of us (JLP) is grateful to the groups at University of Durham, Dept. of Mathe-

matics, and at University of Cambridge, DAMTP for discussions relevant to this work, in particular very useful discussions with A. Kent and A. Taormina are gratefully acknowledged, as is partial support under EEC grant SC1 0394-C(EDB).

References

- [1] K.Ito, J.O. Madsen and J.L. Petersen, NBI-HE-92-42.
- [2] K.Ito and J.O. Madsen, Phys. Lett. **B 283** (1992) 223.
- [3] V.G. Drinfeld and V.V. Sokolov, J. Sov. Math. **30** (1984) 1975; A. Belavin, in “Quantum String Theory” Proceedings in Physics vol. **31** (Springer Verlag, Berlin ,1989); A. Alekseev and S. Shatashvili, Nucl. Phys. **B 323** (1989) 719; M. Bershadsky and H. Ooguri, Commun. Math. Phys. **126** (1989) 429; J. Balog, L. Fehér, P. Forgács, L. O’Raifeartaigh and A. Wipf, Ann. Phys. **203** (1990) 76; A.M. Polyakov, Int. J. Mod. Phys. **A5** (1990) 833; M. Bershadsky, Commun. Math. Phys. **139** (1991)71; F.A. Bais, T.Tjin and P. van Driel, Nucl. Phys. **B 357** (1991) 632; L. Romans, Nucl. Phys. **B 357** (1991) 549. T.G. Khovanova, Theor. Math. Phys. **72** (1987) 899;
- [4] P. Mathieu, Phys. Lett. **B218** (1989) 185.
- [5] K. Miki, Int. J.of Mod. Phys. **A 5** (1990) 1293
- [6] M. Ademollo et al., Phys. Lett. **B 62** (1976) 105; Nucl. Phys. **B 114** (1976) 297.
- [7] S. Matsuda, Phys. Lett. **B 282** (1992) 56.
- [8] A. Sevrin, W. Troost and A. Van Proeyen, Phys. Lett. **B208** (1988) 447.
A. Sevrin, Ph. Spindel, W.Troost and A. Van Proeyen , Nucl. Phys. **B308** (1988) 662; Nucl. Phys. **B311** (1988) 465.
- [9] E.A. Ivanov, S.O. Krivonos and V.M. Leviant, Nucl. Phys **B304** (1988) 601;
Phys. Lett. **B215** (1988) 639.

- [10] V.G. Knizhnik, Theor. Math. Phys. **66** (1986) 68. M. Bershadsky, Phys. Lett. **B174** (1986) 285.
- [11] K. Schoutens, Nucl. Phys. **B314** (1989) 519.
- [12] M. Günaydin, J.L. Petersen, A. Taormina and A. Van Proeyen, Nucl. Phys. **B322** (1989) 402.
- [13] A. Van Proeyen, Class. Quant. Grav. **6** (1989) 1501.
- [14] P. Goddard and A. Schwimmer, Phys. Lett. **B214** (1988) 209.
- [15] J.L. Petersen and A. Taormina, Nucl. Phys. **B331** (1990) 556; Nucl. Phys. **B333** (1990) 833.
- [16] J.L. Petersen and A. Taormina, Nucl. Phys. **B354** (1991) 689. H. Ooguri, J.L. Petersen and A. Taormina, Nucl. Phys. **B368** (1992) 611.
- [17] F. Defever, S. Schrans and K. Thielemans, Phys. Lett. **B212** (1988) 467.
- [18] V.G. Kac, Adv. Math. **26** (1977) 8; Lect. Notes in Math. vol. **676**, p.597 (Springer-Verlag, Berlin, 1978).
- [19] M. Wakimoto, Commun. Math. Phys. **104** (1986) 605; D. Bernard and G. Felder, Commun. Math. Phys. **127** (1990) 145; B. Feigin and E. Frenkel, in “Physics and Mathematics of Strings”, (eds. L. Brink et al., World Scientific, 1990 Singapore).
- [20] T. Eguchi and A. Taormina, Phys. Lett **B196** (1987) 75; *ibid.* **B200** (1988) 315; *ibid.* **B210** (1988) 125.
- [21] V.G. Kac and D.H. Peterson, Adv. in Math. **53** (1984) 125.
- [22] A Kent, and H. Riggs, Phys. Lett. **B198** (1987) 491.

- [23] M. Kato and S. Matsuda, Phys. Lett. **172 B** (1986) 216; Prog. Theor. Phys. **78** (1987) 158; Phys. Lett. **184 B** (1987) 184.